



**COVID-19 TELEHEALTH AND TELEREHABILITATION
REFLECTIONS: IDENTIFYING SOCIOECONOMIC DISPARITIES &
FORMULATING DESIGN RECOMMENDATIONS TO IMPROVE
REMOTE STROKE REHABILITATION**

Frances Watson¹ & Tobi Majekodunmi² Adegboyega Akinsiku³, Dr. Helena Mentis³

¹Department of Computer Science and Electrical Engineering, University of Maryland Baltimore County, 1000 Hilltop Circle, Catonsville, Maryland 2125-

²Department of Mechanical Engineering, University of Maryland Baltimore County, 1000 Hilltop Circle, Catonsville, Maryland 21250

³Department of Information Systems, University of Maryland Baltimore County, 1000 Hilltop Circle, Catonsville, Maryland 21250

Stroke is the leading cause of long-term disability in the United States, and rehabilitation is an important aspect of a stroke survivors' recovery process. However, due to the disruptive nature of the COVID-19 pandemic, health services and rehabilitation for stroke survivors have needed to be achieved remotely. Thus, telehealth and telerehabilitation has become a critical element in providing effective healthcare services. The following study was motivated to further understand the socioeconomic disparities that providers and patients experienced using telerehabilitation and inform the design of future telerehabilitation systems.

We conducted 20 semi-structured interviews that ranged between 45-60 minutes. In light of the COVID-19 pandemic, these interviews took place online via video conferencing platforms (e.g., Cisco Webex). The interviews were with medical rehabilitation specialists (Physiatrists, Occupational Therapists, Physical Therapists, Rehabilitation Psychologists, Speech-Language Pathologists). We qualitatively analyzed interview/transcripts (i.e., open coded), and our analysis revealed several themes. Some of our findings are: (1) caregiver's role in effective telerehabilitation, (2) low health literacy of patients, (3) healthcare is more accessible to a specific population, (4) language barriers impact the efficacy of telerehabilitation, (5) there is a loss of collegiality amongst colleagues, and (6) a virtual telerehabilitation system must be supplemented with physical objects. In the future, we will interview stroke survivors and caregivers and develop an ideal telerehabilitation platform using the design requirements developed from our findings.

We would like to acknowledge the Undergraduate Research Award program, USM LSAMP Research Fellowship program supported by NSF Grant # 1619676, and NSF grant #1552837 for supporting our research.